

PROPER ACTIONS ON TOPOLOGICAL GROUPS: APPLICATIONS TO QUOTIENT SPACES

SERGEY A. ANTONYAN

ABSTRACT. Let X be a Hausdorff topological group and G a locally compact subgroup of X . We show that the natural action of G on X is proper in the sense of R. Palais. This is applied to prove that there exists a closed set $F \subset X$ such that $FG = X$ and the restriction of the quotient projection $X \rightarrow X/G$ to F is a perfect map $F \rightarrow X/G$. This is a key result to prove that many topological properties (among them, paracompactness and normality) are transferred from X to X/G , and some others are transferred from X/G to X . Yet another application leads to the inequality $\dim X \leq \dim X/G + \dim G$ for every paracompact group X and its locally compact subgroup G .

1. INTRODUCTION

By a G -space we mean a completely regular Hausdorff space together with a fixed continuous action of a Hausdorff topological group G on it.

The notion of a proper G -space was introduced in 1961 by R. Palais [24] with the purpose to extend a substantial portion of the theory of compact Lie group actions to the case of noncompact ones.

A G -space X is called proper (in the sense of Palais [24, Definition 1.2.2]), if each point of X has a, so called, *small* neighborhood, i.e., a neighborhood V such that for every point of X there is a neighborhood U with the property that the set $\langle U, V \rangle = \{g \in G \mid gU \cap V \neq \emptyset\}$ has compact closure in G . Clearly, if G is compact, then every G -space is proper.

Many important problems in the theory of proper actions are conjugated (see [19], [1], [2], [8], [7]) to the following major open problem:

Conjecture 1. *Let G be a locally compact group. Then the orbit space X/G of any paracompact proper G -space X is paracompact.*

This conjecture is open even if X is metrizable; in this case it is equivalent (see [8]) to the following old problem going back to R. Palais [24]:

Conjecture 2. *Let G be a locally compact group and X a metrizable proper G -space. Then the topology of X is metrizable by a G -invariant metric.*

Due to Palais [24], it is known that Conjecture 2 is true for separable metrizable proper G -spaces provided the acting group G is Lie. Other particular cases are discussed in [21] and [8].

2000 *Mathematics Subject Classification.* 22A05; 22F05; 54H11; 54H15; 54F45.

Key words and phrases. Proper G -space; orbit space; locally compact group; paracompact space; dimension.

The author was supported in part by grant IN-102608 from PAPIIT (UNAM).

In this paper we prove Conjecture 1 in an important particular case, namely, when X is a topological group endowed with the natural action of its locally compact subgroup G . (see Corollary 1.5). We first show that X is a proper G -space and then we establish a more general result (Theorem 1.2) which has many interesting applications in the theory of topological groups.

Below all topological groups are assumed to satisfy the Hausdorff separation axiom.

Theorem 1.1. *Let X be a topological group and G a locally compact subgroup of X . Then the action of G on X given by the formula $g * x = xg^{-1}$, $g \in G$, $x \in X$, is proper.*

Recall that a subset S of a proper G -space X is called G -fundamental, or just fundamental, if S is a small set and the saturation $G(S) = \{gs \mid g \in G, s \in S\}$ coincides with X .

Here is the key result of the paper:

Theorem 1.2. *Let X be a topological group and G a locally compact subgroup of X . Then there exists a closed G -fundamental set in X .*

It is easy to prove (see [2, Proposition 1.4]) that in every proper G -space X , the restriction of the orbit map $p : X \rightarrow X/G$ to any closed small set is perfect (i.e., is closed and has compact fibers). In combination with Theorem 1.2 this yields:

Corollary 1.3. *Let X be a topological group, G a locally compact subgroup of X , and X/G the quotient space of all right lateral classes $xG = \{xg \mid g \in G\}$, $x \in X$. Then there exists a closed subset $F \subset X$ such that the restriction $p|_F : F \rightarrow X/G$ is a perfect surjective map.*

This fact has the following immediate corollary about transfer of properties from X to X/G :

Corollary 1.4. *Let $\mathcal{P}$ be a topological property stable under perfect maps and also inherited by closed subsets. Assume that X is a topological group with the property $\mathcal{P}$ and let G be a locally compact subgroup of X . Then the quotient space X/G also has the property $\mathcal{P}$.*

Among such properties $\mathcal{P}$ we single out just some of those which provide new results in Corollary 1.4; these are: paracompactness, countable paracompactness, weak paracompactness, normality, perfect normality, Čech-completeness, being a k -space (see [16, §5.1, §5.2, §5.3, §1.5, §3.9, §3.3]) and stratifiability (see [14]). Thus, we get the following positive solution of Conjecture 1 in an important special case:

Corollary 1.5. *Let X be a paracompact topological group and G a locally compact subgroup of X . Then the quotient space X/G is paracompact.*

In this connection it is in order to recall the following remarkable result of A. V. Arhangel'skii [9]: every topological group is the quotient of a paracompact zero-dimensional group. Hence, the local compactness of G is essential in Corollary 1.5.

We recall that a locally compact group is called *almost connected*, if its space of connected components is compact.

Corollary 1.6. *Let $\mathcal{P}$ be a topological property stable under open perfect maps and also inherited by closed subsets. Assume that X is a paracompact group with the property $\mathcal{P}$ and let G be an almost connected subgroup of X . Then the quotient space X/G also has the property $\mathcal{P}$.*

Among properties stable under open perfect maps and also inherited by closed subsets we highlight strong paracompactness and realcompactness (see [16, Exercises 5.3.C(c), 5.3H(d), and Theorem 3.11.4 and Exercises 3.11.G]). Thus, we get the following

Corollary 1.7. *Let X be a strongly paracompact (resp., paracompact and realcompact) topological group and G an almost connected subgroup of X . Then the quotient space X/G is strongly paracompact (resp., paracompact and realcompact).*

We note that the converse of the first statement in this corollary is not true. Namely, it is known that the Baire space $B(\aleph_1)$ of weight $\aleph_1$ (which is, in fact, a commutative topological group) is strongly paracompact and its product with the additive group $\mathbb{R}$ of the reals is not (see [16, Exercises 5.3F(a) and 5.3F(b)]). Hence, the direct product $X = \mathbb{R} \times B(\aleph_1)$ and its subgroup $G = \mathbb{R} \times \{0\}$ provide the desired counterexample, answering negatively a question from [11]. Furthermore, [12, Open Problem 3.2.1, p.151] asks whether every locally strongly paracompact group is strongly paracompact? The same group $\mathbb{R} \times B(\aleph_1)$ provides a negative answer to this question too. Indeed, since the product of a compact space and a strongly paracompact space is strongly paracompact (see [16, Exercise 5.3H(a)]), we infer that $\mathbb{R} \times B(\aleph_1)$ is locally strongly paracompact.

Combining our Corollary 1.5 with a result of Abels [1, Main Theorem], we obtain the following

Corollary 1.8. *Let X be a paracompact group, G an almost connected subgroup of X , and K a maximal compact subgroup of G . Then there exists a K -invariant subset $S \subset X$ such that X is K -homeomorphic to the product $G/K \times S$. In particular, X is homeomorphic to $\mathbb{R}^n \times S$ for some $n \geq 1$.*

In [10] A. V. Arhangel'skii has studied properties which are transferred in the opposite direction, i.e., from X/G to X . The next corollary is a unified result of this sort which implies many of those in [10] as well as provides some new ones.

Corollary 1.9. *Let $\mathcal{P}$ be a topological property invariant and inverse invariant of perfect maps, and also stable under multiplication by a locally compact group. Assume that X is a topological group and let G be a locally compact subgroup of X such that the quotient space X/G has the property $\mathcal{P}$. Then the group X also has the property $\mathcal{P}$.*

Among such properties we mention just some: paracompactness, being a k -space, Čech-completeness (see [16, §5.1, §3.3, §3.9]). That paracompactness is stable under multiplication by a locally compact group follows from a result of Morita [22] since every locally compact group is paracompact (even, strongly paracompact [12, Theorem 3.1.1]).

Corollary 1.10. *Let $\mathcal{P}$ be a topological property invariant and inverse invariant of open perfect maps, and also stable under multiplication by a locally compact group. Assume that X is a paracompact group and let G be an almost connected subgroup*

of X such that the quotient space X/G has the property $\mathcal{P}$. Then the group X also has the property $\mathcal{P}$.

Among such properties we highlight realcompactness (see [16, Theorem 3.11.14 and Exercise 3.11.G, and also take into account that every locally compact group is realcompact]).

Coorollary 1.5 is further applied to prove the following Hurewicz type formula:

Theorem 1.11. *Let G be a locally compact subgroup of a paracompact topological group X . Then*

$$\dim X \leq \dim X/G + \dim G.$$

Remark 1.12 ([25]). *If in this theorem X is a locally compact group then, in fact, the equality holds:*

$$\dim X = \dim X/G + \dim G.$$

All the proofs are given in section 3.

2. PRELIMINARIES

Throughout the paper, unless otherwise is stated, by a *group* we shall mean a topological group G satisfying the Hausdorff separation axiom; by e we shall denote the unity of G .

All topological spaces are assumed to be Tychonoff (= completely regular and Hausdorff). The basic ideas and facts of the theory of G -spaces or topological transformation groups can be found in G. Bredon [15] and in R. Palais [23]. Our basic reference on proper group actions is Palais' article [24]. Other good sources are [21], [1] and [2].

For the convenience of the reader we recall, however, some more special definitions and facts below.

By a G -space we mean a topological space X together with a fixed continuous action $G \times X \rightarrow X$ of a topological group G on X . By gx we shall denote the image of the pair $(g, x) \in G \times X$ under the action.

If Y is another G -space, a continuous map $f : X \rightarrow Y$ is called a G -map or an equivariant map, if $f(gx) = gf(x)$ for every $x \in X$ and $g \in G$.

If X is a G -space, then for a subset $S \subset X$ and for a subgroup $H \subset G$, the H -hull (or H -saturation) of S is defined as follows: $H(S) = \{hs \mid h \in H, s \in S\}$. If S is the one point set $\{x\}$, then the G -hull $G(\{x\})$ usually is denoted by $G(x)$ and called the orbit of x . The orbit space X/G is always considered in its quotient topology.

A subset $S \subset X$ is called H -invariant if it coincides with its H -hull, i.e., $S = H(S)$. By an invariant set we shall mean a G -invariant set.

For any $x \in X$, the subgroup $G_x = \{g \in G \mid gx = x\}$ is called the stabilizer (or stationary subgroup) at x .

A compatible metric ρ on a metrizable G -space X is called invariant or G -invariant, if $\rho(gx, gy) = \rho(x, y)$ for all $g \in G$ and $x, y \in X$. If ρ is a G -invariant metric on any G -space X , then it is easy to verify that the formula

$$\tilde{\rho}(G(x), G(y)) = \inf\{\rho(x', y') \mid x' \in G(x), y' \in G(y)\}$$

defines a pseudometric $\tilde{\rho}$, compatible with the quotient topology of X/G . If, in addition, X is a proper G -space then $\tilde{\rho}$ is, in fact, a metric on X/G [24, Theorem 4.3.4].

For a closed subgroup $H \subset G$, by G/H we will denote the G -space of cosets $\{gH \mid g \in G\}$ under the action induced by left translations.

A locally compact group G is called *almost connected*, if the space of connected components of G is compact. Such a group has a maximal compact subgroup K , i.e., every compact subgroup of G is conjugate to a subgroup of K [1, Theorem A.5]. The corresponding classical theorem on Lie groups can be found in [20, Ch. XV, Theorem 3.1].

In 1961 Palais [24] introduced the very important concept of a *proper action* of an arbitrary locally compact group G and extends a substantial part of the theory of compact Lie transformation groups to noncompact ones.

Let X be a G -space. Two subsets U and V in X are called thin relative to each other [24, Definition 1.1.1], if the set $\langle U, V \rangle = \{g \in G \mid gU \cap V \neq \emptyset\}$, called *the transporter* from U to V , has a compact closure in G . A subset U of a G -space X is called *G -small*, or just *small*, if every point in X has a neighborhood thin relative to U . A G -space X is called *proper* (in the sense of Palais), if every point in X has a small neighborhood.

Clearly, if G is compact, then every G -space is proper. Furthermore, if G acts properly on a compact space, then G has to be compact as well. If G is discrete and X is locally compact, the notion of a proper action is the same as the classical notion of a *properly discontinuous* action. When $G = \mathbb{R}$, the additive group of the reals, proper G -spaces are precisely the *dispersive* dynamical systems with regular orbit space (see [13, Ch. IV]).

Important examples of proper G -spaces are the coset spaces G/H with H a compact subgroup of a locally compact group G . Other interesting examples the reader can find in [1], [2], [5], [6] and [21].

In what follows we shall need also the definition of a twisted product $G \times_K S$, where K is a closed subgroup of G , and S a K -space. $G \times_K S$ is the orbit space of the K -space $G \times S$ on which K acts by the rule: $k(g, s) = (gk^{-1}, ks)$. Furthermore, there is a natural action of G on $G \times_K S$ given by $g'[g, s] = [g'g, s]$, where $g' \in G$ and $[g, s]$ denotes the K -orbit of the point (g, s) in $G \times S$. We shall identify S , by means of the K -equivariant embedding $s \mapsto [e, s]$, $s \in S$, with the K -invariant subset $\{[e, s] \mid s \in S\}$ of $G \times_K S$. This K -equivariant embedding $S \hookrightarrow G \times_K S$ induces a homeomorphism of the K -orbit space S/K onto the G -orbit space $(G \times_K S)/G$ (see [15, Ch. II, Proposition 3.3]).

The twisted products are of a particular interest in the theory of transformation groups (see [15, Ch. II, § 2]). It turns out that every G -space locally is a twisted product. For a more precise formulation we need to recall the following well known notion of a slice (see [24, p. 305]):

Definition 2.1. *Let X be a G -space and K a closed subgroup of G . A K -invariant subset $S \subset X$ is called a K -kernel if there is a G -equivariant map $f : G(S) \rightarrow G/K$, called the *slicing map*, such that $S = f^{-1}(eK)$. The saturation $G(S)$ is called a *tubular set* and the subgroup K will be referred as the *slicing subgroup*.*

If in addition $G(S)$ is open in X then we shall call S a K -slice in X .

If $G(S) = X$ then S is called a global K -slice of X .

It turns out that each tubular set with a compact slicing subgroup is a twisted product. The tubular neighborhood $G(S)$ is G -homeomorphic to the twisted product $G \times_K S$; namely the map $\xi : G \times_K S \rightarrow G(S)$ defined by $\xi([g, s]) = gs$ is a G -homeomorphism (see [15, Ch. II, Theorem 4.2]). In what follows we will use this fact without a specific reference.

One of the most powerful results in the theory of topological transformation groups states (see [24, Proposition 2.3.1]) that, if X is a proper G -space with G a Lie group, then for any point $x \in X$, there exists a G_x -slice S in X such that $x \in S$. In general, when G is not a Lie group, it is no longer true that a slice exists at each point of X (see [4]). Generalizing the case of Lie group actions, in [7] the following approximate version of Palais' Slice Theorem [24, Proposition 2.3.1] for non-Lie group actions was proved, which plays a key role in the proof of Theorem 1.11:

Theorem 2.2 (Approximate Slice Theorem [7]). *Let G be a locally compact group, X a proper G -space and $x \in X$. Then for any neighborhood O of x in X , there exist a compact large subgroup $K \subset G$ with $G_x \subset K$, and a K -slice S such that $x \in S \subset O$.*

Thus, every proper G -space is covered by invariant open subsets each of which is a twisted product.

Note that for G a compact group this theorem was proved in [4]. A version of it, without requiring K to be a large subgroup, was obtained in [2].

We recall that here a closed subgroup $H \subset G$ is called *large*, if there exists a closed normal subgroup $N \subset G$ such that $N \subset H$ and G/N is a Lie group.

We refer to [7] for further characterizations of large subgroups.

On any group G one can define two natural (but equivalent) actions of G given by the formulas

$$g \cdot x = gx, \quad \text{and} \quad g * x = xg^{-1},$$

respectively, where in the right parts the group operations are used with $g, x \in G$.

Throughout we shall consider the second action only.

By $U(G)$ we shall denote the Banach space of all right uniformly continuous bounded functions $f : G \rightarrow \mathbb{R}$ endowed with the supremum norm. Recall that f is called right uniformly continuous, if for every $\varepsilon > 0$ there exists a neighborhood O of the unity in G such that $|f(y) - f(x)| < \varepsilon$ whenever $yx^{-1} \in O$.

We shall consider the induced action of G on $U(G)$, i.e.,

$$(gf)(x) = f(xg), \quad \text{for all } g, x \in G.$$

It is easy to check that this action is continuous, linear and isometric (see e.g., [3, Proposition 7]).

Proposition 2.3. *Let G be a group. Then for every $f \in U(G)$, the map $f_* : G \rightarrow U(G)$ defined by $f_*(x)(g) = f(xg^{-1})$, $x, g \in G$, is a right uniformly continuous G -map.*

Proof. A simple verification. □

3. PROOFS

Proof of Theorem 1.1. Choose a neighborhood U of the identity in X such that $U = U^{-1}$ and $U^3 \cap G$ has a compact closure in G . We claim that for every $x \in X$ the neighborhood xU is G -small. Indeed, let $y \in X$ be any point. Two cases are possible.

Case 1. Assume that $y \in xU^2G$. Then $y = xu_1u_2h$ with $u_1, u_2 \in U$ and $h \in G$. We claim that xU^2h is a neighborhood of y thin relative to xU . Indeed, if $g \in \langle xU, xU^2h \rangle$, then $g^{-1}h^{-1} \in U^3 \cap G$. Since $U^3 \cap G$ has a compact closure, we see that so is the set $(U^3 \cap G)h$ to which belongs g^{-1} . This yields that $\langle xU, xU^2h \rangle$ is contained in $h^{-1}(U^3 \cap G)^{-1}$, which also has a compact closure. Hence the transporter $\langle xU, xU^2h \rangle$ has a compact closure, as required.

Case 2. Assume that $y \notin xU^2G$. In this case $y \notin \overline{xUG}$. Indeed, if $y \in \overline{xUG}$ then the neighborhood $xUx^{-1}y$ of y should meet the set xUG . Then $xux^{-1}y = xvh$ for suitable elements $u, v \in U$ and $h \in G$. Then $y = xu^{-1}vh \in xU^2G$, a contradiction.

Hence the open set $V = X \setminus \overline{xUG}$ is a G -invariant neighborhood of y , and V is thin relative to xU because the transporter $\langle xU, V \rangle$ is empty in this case. This completes the proof.

Proposition 3.1. *Let X be a group and G a locally compact subgroup of X . Then there exists a locally finite covering of X consisting of G -invariant open sets of the form S_iG , where each S_i is an open G -small subset of X .*

Proof. By Theorem 1.1, X is a proper G -space, and hence, one can choose a G -small neighborhood U of the unity in X . By virtue of Markov's theorem [12, Theorem 3.3.9], there exists a right uniformly continuous function $f : X \rightarrow [0, 1]$ such that

$$(3.1) \quad f(e) = 0 \quad \text{and} \quad f^{-1}([0, 1)) \subset U.$$

Then, by Proposition 2.3, f induces an X -equivariant map $f_* : X \rightarrow U(X)$ defined by the rule:

$$f_*(x)(g) = f(xg^{-1}), \quad x, g \in X.$$

Denote by Z the image $f_*(X)$. Clearly, Z is the X -orbit of the point $f_*(e)$ in the X -space $U(X)$, and the metric of $U(X)$ induces an X -invariant metric on Z . We claim that

$$(3.2) \quad f_*^{-1}(\Gamma_{x,V}) \subset x^{-1}U, \quad \text{for every } x \in X,$$

where $V = [0, 1)$ and $\Gamma_{x,V}$ is the open subset $\{\varphi \in U(X) \mid \varphi(x) \in V\}$ of U .

First we observe that $\Gamma_{x,V} = x^{-1}\Gamma_{e,V}$ and

$$f_*^{-1}(\Gamma_{e,V}) \subset f^{-1}(V).$$

Then (3.2) follows from (3.1) and the X -equivariance of f_* .

Besides, since $f_*(x) \in \Gamma_{x,V}$ for every $x \in X$, we see that the sets $\Gamma_{x,V}$, $x \in X$, constitute a covering of Z .

From now on we restrict ourselves only by the induced actions of the subgroup G , i.e., we will consider X and Z just as G -spaces.

Now, it follows from (3.2) and from the G -equivariance of f_* that

$$(3.3) \quad f_*^{-1}(G(\Gamma_{x,V})) \subset x^{-1}UG, \quad \text{for every } x \in X.$$

Since $f_* : X \rightarrow Z$ is G -equivariant, it induces a continuous map $\tilde{f}_*$ of the G -orbit spaces, i.e., we have the following commutative diagram:

$$\begin{array}{ccc} X & \xrightarrow{f_*} & Z \\ \downarrow p & & \downarrow q \\ X/G & \xrightarrow{\tilde{f}_*} & Z/G \end{array}$$

where p and q are the G -orbit maps.

It follows from (3.3) that

$$(3.4) \quad \tilde{f}_*^{-1}(q(\Gamma_{x,V})) \subset p(x^{-1}U), \quad \text{for every } x \in X.$$

Thus, the open covering $\{p(xU) \mid x \in X\}$ of the G -orbit space X/G is refined by the open covering $\{\tilde{f}_*^{-1}(q(\Gamma_{x,V})) \mid x \in X\}$.

Since the metric of Z is G -invariant, the orbit space Z/G is pseudometrizable (see Preliminaries). Hence the open covering $\{q(\Gamma_{x,V}) \mid x \in X\}$ of Z/G admits an open locally finite refinement, say $\{W_i \mid i \in \mathcal{I}\}$ (see [16, Theorem 4.4.1 and Remark 4.4.2]). Then, clearly, $\{\tilde{f}_*^{-1}(W_i) \mid i \in \mathcal{I}\}$ is an open locally finite refinement of $\{xUG \mid x \in X\}$ consisting of G -invariant sets. It then follows that each set $p^{-1}(\tilde{f}_*^{-1}(W_i))$ is contained in some set xUG , $x \in X$, which yields that

$$p^{-1}(\tilde{f}_*^{-1}(W_i)) = S_i G,$$

where $S_i = p^{-1}(\tilde{f}_*^{-1}(W_i)) \cap xU$.

Now, S_i , being a subset of the G -small set xU , is itself G -small (see the proof of Theorem 1.1). This completes the proof. $\square$

Proof of Theorem 1.2. Let $\{S_i G \mid i \in \mathcal{I}\}$ be the locally finite open covering of X from Proposition 3.1. Then the union $S = \bigcup_{i \in \mathcal{I}} S_i$ is a G -small set (see e.g., [2, Proposition 1.2(d)]). On the other hand,

$$SG = \left(\bigcup_{i \in \mathcal{I}} S_i \right) G = \bigcup_{i \in \mathcal{I}} S_i G = X,$$

yielding that S is a G -fundamental subset of X . Since the closure of a G -small set is G -small (see e.g., [2, Proposition 1.2(b)]), the closure $\overline{S}$ is the desired closed G -fundamental set. This completes the proof.

Proof of Corollary 1.6. Since G is almost connected, it has a maximal compact subgroup K (see Preliminaries). Since by Corollary 1.5, the quotient X/G is para-compact, due to a result of Abels [1, Main Theorem], X admits a global K -slice S , and hence, it is G -homeomorphic to the twisted product $G \times_K S$ (see Preliminaries). Since the group K is compact, it then follows that the K -orbit map $G \times S \rightarrow G \times_K S \cong_G X$ is open and perfect. This yields immediately that the restriction $p|_S : S \rightarrow X/G$ of the G -orbit map $p : X \rightarrow X/G$ is an open and perfect surjection. Now the result follows.

Proof of Corollary 1.9. By virtue of Corollary 1.3, X admits a closed subset $F \subset X$ such that the restriction of the quotient projection $p : X \rightarrow X/G$ to F is a

perfect surjection $p|_F : F \rightarrow X/G$. It then follows from the hypothesis that F has the property $\mathcal{P}$. Since a locally compact group is paracompact (even, strongly paracompact [12, Theorem 3.1.1]), then again by the hypothesis, the product $G \times F$ has the property $\mathcal{P}$. Since F is a closed G -small set, the action map $G \times F \rightarrow X$ is perfect (see Abels [2, Proposition 1.4]), and since $FG = X$ we see that the map $G \times F \rightarrow X$ is surjective. Then X has the property $\mathcal{P}$ by the hypothesis.

Proof of Corollary 1.10. It is quite similar to the proof of Corollary 1.6.

Proof of Theorem 1.11. Consider three cases.

Casa 1. Assume that G is almost connected. In this case G has a maximal compact subgroup, say, K (see Preliminaries). Since, by Corollary 1.5, the quotient X/G is paracompact, one can apply a theorem of H. Abels [1, Main Theorem], according to which X admits a global K -slice, say S , and hence, X is G -homeomorphic to the twisted product $G \times_K S$. By another result of H. Abels [1, Theorem 2.1], X is homeomorphic (moreover, K -equivariantly homeomorphic) to the product $G/K \times S$.

Since G is locally compact and paracompact and the natural quotient map $G \rightarrow G/K$ is open and closed, we infer that G/K is also locally compact and paracompact. Further, since S is paracompact, according to a theorem of Morita [22] one has:

$$(3.5) \quad \dim(G/K \times S) \leq \dim G/K + \dim S.$$

Since K is compact, according to a result of V. V. Filippov [18] one has the inequality:

$$(3.6) \quad \dim S \leq \dim S/K + \dim q$$

where $q : S \rightarrow S/K$ is the K -orbit projection and $\dim q = \sup \{\dim q^{-1}(a) \mid a \in S/K\}$. Further, since K acts freely on S , we see that $\dim q = \dim K$.

Consequently, combining (3.5) and (3.6) one obtains:

$$(3.7) \quad \dim(G/K \times S) \leq \dim G/K + \dim K + \dim S/K.$$

Next, since $X/G \cong (G \times_K S)/G \cong S/K$ (see Preliminaries) and $\dim G/K + \dim K = \dim G$ (see Remark 1.12), it then follows from (3.7) that

$$\dim(G/K \times S) \leq \dim X/G + \dim G,$$

as required.

Case 2. Let G be totally disconnected. Then G is zero-dimensional (see [17, Theorem 1.4.5]). Hence, one has to prove that

$$\dim X \leq \dim X/G.$$

By virtue of Approximate Slice Theorem (see Theorem 2.2), X is covered by a family of open tubular sets U_i . Assume that $f_i : U_i \rightarrow G/K_i$ is the corresponding slicing map, where each K_i is a compact large subgroup of G .

Let $p : X \rightarrow X/G$ be the quotient map. Since, by Corollary 1.5, X/G is paracompact the open covering $\{p(U_i)\}$ of X/G has a locally finite closed shrinking $\{F_i\}$, i.e., $F_i \subset p(U_i)$ and $\bigcup F_i = X/G$ (see [16, Remark 5.1.7]). Then, putting $\Phi_i = p^{-1}(F_i)$, we get a locally finite closed shrinking $\{\Phi_i\}$ of $\{U_i\}$ consisting of G -invariant sets.

Thus, X is covered by a locally finite family of closed tubular sets Φ_i . Consequently, by virtue of the locally finite sum theorem (see [17, Theorem 3.1.10]), it suffices to show that $\dim \Phi \leq \dim X/G$, for every member Φ of the covering $\{\Phi_i\}$.

Let $\varphi : \Phi \rightarrow G/K$ be the slicing map corresponding to a tubular set $\Phi \in \{\Phi_i\}$, where K is a compact large subgroup of G (of course, depending upon Φ). Since K is a large subgroup, the quotient G/K is locally connected, in fact, it is a manifold [7, Proposition 3.2]. On the other hand G/K is totally disconnected, and hence, G/K should be a discrete space. This yields that, if we put $Q = f^{-1}(eK)$, then each gQ is closed and open in Φ , and Φ is the disjoint union of the sets gQ one g out of every coset in G/K . In other words, Φ is just homeomorphic to the product $G/K \times Q$.

Next, since G/K is locally compact and paracompact, and Q is paracompact (being a closed subset of the paracompact space X), according to a theorem of Morita [22] one has:

$$(3.8) \quad \dim \Phi = \dim (G/K \times Q) \leq \dim G/K + \dim Q.$$

Since K is compact, according to a result of V. V. Filippov [18] one has the inequality:

$$(3.9) \quad \dim Q \leq \dim Q/K + \dim r$$

where $r : Q \rightarrow Q/K$ is the K -orbit projection. Further, since K acts freely on Q , we see that $\dim r = \dim K \leq \dim G = 0$.

Consequently, combining (3.8) and (3.9), and taking into account that $\dim K = 0 = \dim G/K$ (this follows, for instance, from $\dim G = 0$ and Remark 1.12), one obtains:

$$(3.10) \quad \dim \Phi \leq \dim Q/K.$$

Further, since Φ is a tubular set with the K -slice Q , one has $\Phi/G \cong Q/K$ (see Preliminaries). Then it follows from (3.10) that

$$(3.11) \quad \dim \Phi \leq \dim \Phi/G.$$

But Φ/G is a closed subset of X/G , and since X/G is normal, due to monotonicity of the dimension (see [17, Theorem 3.1.4]), one has $\dim \Phi/G \leq \dim X/G$. This, together with (3.11), now yields that

$$\dim \Phi \leq \dim X/G,$$

as required.

Case 3. Let G be arbitrary. By case 1, we have:

$$(3.12) \quad \dim X \leq \dim X/G_0 + \dim G_0,$$

where G_0 is the identity component of G . Since G/G_0 is a totally disconnected locally compact group which acts naturally on the G_0 -orbit space X/G_0 , then according to case 2, one has:

$$(3.13) \quad \dim X/G_0 \leq \dim \frac{X/G_0}{G/G_0}.$$

Now, since $\frac{X/G_0}{G/G_0} \cong X/G$ and $\dim G_0 \leq \dim G$, it follows from (3.12) and (3.13) that

$$\dim X \leq \dim X/G + \dim G.$$

This completes the proof.

REFERENCES

1. H. Abels, *Parallelizability of proper actions, global K -slices and maximal compact subgroups*, Math. Ann. **212** (1974), 1–19.
2. H. Abels, *A universal proper G -space*, Math. Z. **159** (1978), 143–158.
3. S. A. Antonian, *Equivariant embeddings into G -AR's*, Glasnik Matematički **22 (42)** (1987), 503–533.
4. S. A. Antonyan, *Existence of a slice for arbitrary compact transformation groups*, Matematicheskie Zametki **56:5** (1994) 3–9: English transl. in: Math. Notes **56 (5-6)** (1994) 101–1104.
5. S. A. Antonyan, *The Banach-Mazur compacta are absolute retracts*, Bull. Acad. Polon. Sci. Ser. Math. **46, no. 2** (1998) 113–119.
6. S. A. Antonyan, *Extensorial properties of orbit spaces of proper group actions*, Topol. Appl. **98** (1999), 35–46.
7. S. A. Antonyan, *Orbit spaces and unions of equivariant absolute neighborhood extensors*, Topology Appl. **146-147** (2005) 289–315.
8. S. A. Antonyan and S. de Neymet, *Invariant pseudometrics on Palais proper G -spaces*, Acta Math. Hung. **98 (1-2)** (2003), 41–51.
9. A. V. Arhangel'skii, *Classes of topological groups*, Russian Math. Surveys **36, no. 3** (1981), 151–174.
10. A. V. Arhangel'skii, *Quotients with respect to locally compact subgroups*, Houston J. Math. **31, no. 1** (2005), 215–226.
11. A. V. Arhangel'skii and V. V. Uspenskij, *Topological groups: local versus global*, Applied General Topology **7, no. 1** (2006), 67–72.
12. A. Arhangel'skii and M. Tkachenko, *Topological Groups and Related Structures*, Atlantis Press/World Scientific, Amsterdam-Paris, 2008.
13. N. P. Bhatia, G. P. Szegö, *Stability theory of Dynamical Systems*, Springer Verlag, Berlin, 1970.
14. C. J. R. Borges, *On stratifiable spaces*, Pacific J. Math. **17, no. 1** (1966), 1–16.
15. G. Bredon, *Introduction to compact transformation groups*, Academic Press, 1972.
16. R. Engelking, *General Topology*, PWN-Pol. Sci. Publ., Warsaw, 1977.
17. R. Engelking, *Dimension Theory*, PWN-Pol. Sci. Publ., Warsaw, 1978.
18. V. V. Filippov, *Dimensionality of spaces with the action of a bicomact group*, Math. Notes, **25, no. 3** (1979), 171–174.
19. O. Hájek, *Parallelizability revisited*, Proc. Amer. Math. Soc., **27, no. 1** (1971), 77–84.
20. G. Hochschild, *The structure of Lie groups*, Holden-Day Inc., San Francisco, 1965.
21. J. L. Koszul, *Lectures on groups of transformations*, Tata Institute of Fundamental Research, Bombay, 1965.
22. K. Morita, *On the Dimension of Product Spaces*, American Journal of Mathematics, Vol. 75, No. 2 (1953), pp. 205–223.
23. R. Palais, *The classification of G -spaces*, Memoirs of the AMS, **36** (1960).
24. R. Palais, *On the existence of slices for actions of non-compact Lie groups*, Ann. of Math. **73**, (1961), 295–323.
25. E. G. Skljarenko, *On the topological structure of locally bicomact groups and their quotient spaces*, Amer. Math. Soc. Transl., Ser. 2, **39** (1964), 57–82.

DEPARTAMENTO DE MATEMÁTICAS, FACULTAD DE CIENCIAS, UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, 04510 MÉXICO DISTRITO FEDERAL, MÉXICO.

E-mail address: antonyan@unam.mx